

TECHNICAL DATA SHEET

Sabouraud Dextrose Broth (Harmonized)

Principle

Sabouraud dextrose broth is used for cultivation of yeasts, molds and aciduric bacteria. It is recommended by USP as a medium for microbial limit testing of pharmaceutical products and raw material used in pharmaceutical industries. The medium is prepared in accordance with the harmonized principles of USP. Medium is consisting of meat and casein peptone (1:1) and dextrose monohydrate. The meat and casein peptone provide carbonaceous, nitrogenous compounds, long chain amino acids, vitamins and other essential growth nutrients in addition to that the dextrose monohydrate serves as energy source. The high concentration of dextrose and low pH of medium favor the growth of yeasts and molds and inhibit other contaminating bacteria from pharmaceutical and clinical specimens.

Use: For selection, isolation & cultivation of yeasts, molds and aciduric microorganisms.

Contents*

Ingredients	Gram/Litre
Meat and Casein Peptone	10.000
Dextrose Monohydrate	20.000
pH at 25°C	5.6 ±0.2

* Formula adjusted for optimum performance and parameters

Directions: Dissolve 30.0 grams in 1000 ml distilled water. Sterilize by autoclaving at 15 lbs. pressure (121°C) for 15 min, cool it to 42-45 °C and inoculate test sample aseptically.

Specimens types analyzed

Pharmaceutical samples, clinical samples, etc.

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Precautions to be taken

These microbial media are intended for the in-vitro use only. All the handling, experiments, storage, and discarding should be performed with the help of skilled and knowledgeable technicians and as per the established guidelines. The material should be disposed only after proper sterilization by autoclaving. Please go through the MSDS of the media to avoid any accidents or in emergency.

Performance and Evaluation

The expected performance of the medium is liable to use as per the direction on the label when stored at optimum conditions and within expiry date.

Quality Control

Appearance	Light beige colored free flowing, homogeneous powder.
Reaction of 3.0% solution	5.6 $\pm$ 0.2 at 25 °C.
pH	5.40- 5.80 at 25 °C.
Color and clarity of ready medium	Light to medium amber, very slightly opalescent clear solution.
Growth Promotion properties	Best at $\leq$ 100 CFU at 22 - 25 °C for 3-5 days
Indicative properties	Optimum at $\leq$ 100 CFU at 22 - 25 °C for 3-5 days.

Different Microbial Response: Cultural characteristics observed after incubation at 25 $\pm$ 2°C for 3-5 days.

Organism	ATCC	Inoculum (CFU)	Growth
<i>Candida albicans</i>	10231	50-100	Luxuriant
<i>Aspergillus brasiliensis</i>	16404	50-100	Luxuriant
<i>Saccharomyces cerevisiae</i>	9763	50-100	Luxuriant

Storage and Shelf Life: The product is highly hygroscopic; keep the container tightly closed at all times and store it properly as per the conditions mentioned on the label. The declared expiry is valid only when stored as per the conditions mentioned on the label. Note: Sterilize media immediately after reconstitution.

This document has been produced electronically and it is valid without signature.

www.oxfordlabchem.com

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Disposal: To avoid the contamination or propagation of any hazardous microbes the used, unusable or modified preparation of this product must be disposed after autoclaving after completion of task.

Reference

1. Atlas, R. M. (2005). Handbook of media for environmental microbiology. CRC press.
2. British Pharmacopoeia, 2016, The Stationery office British Pharmacopoeia.
3. Difco Manual (1998). 11th Edition. Difco Laboratories., Division of Becton Dickinson and Company, Sparks, Maryland, USA.
4. European Pharmacopoeia, 2017, European Dept. for the quality of Medicines.
5. Isenberg, H.D. Clinical Microbiology Procedures Handbook, 2nd Edition.
6. Japanese Pharmacopoeia, 2016.
7. The United States Pharmacopoeia, (2014), The United States Pharmacopeial Convention. 12601 Twinbrook Parkway, Rockville, MD 20852.

Disclaimer:

The information contained herein in good faith but makes no representations as to its comprehensiveness or accuracy. This document is intended only as a guide to the appropriate precautionary handling of the material by a properly trained person using this product. Individuals receiving the information must exercise their independent judgment in determining its appropriateness for a particular purpose.

Oxford Lab Fine Chem LLP makes no representations or warranties, either express or implied, including without limitation any warranties of merchantability, fitness for a particular purpose with respect to the information set forth herein or the product to which the information refers. Accordingly, Oxford Lab Fine Chem LLP will not be responsible for damages resulting from use of or reliance upon this information.

This document has been produced electronically and it is valid without signature.

www.oxfordlabchem.com