

TECHNICAL DATA SHEET

Tryptone Glucose Yeast Extract Broth (Standard Method Broth)

Principle

Tryptone glucose yeast extract agar is simple media used for enumeration and cultivation of bacteria from potable water, waste water, air and food and dairy products. It is composed of tryptone, yeast extract, glucose and agar. Tryptone and yeast extract provide nitrogen, carbon, vitamins and other essential growth factors. Glucose is carbon source. Agar is solidifying agent.

Use: For enumeration of bacteria in water, air, milk and dairy products.

Contents*

Ingredients	Gram/Litre
Tryptone	10.00
Yeast Extract	1.00
Glucose	5.00
Dipotassium Hydrogen Phosphate	1.25
pH at 25°C	6.8 ±0.2

*** Formula adjusted for optimum performance and parameters**

Directions: Dissolve 17.25 grams in 1000 ml distilled water. Boil to dissolve the medium completely and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 min, cool it to 42-45 °C and inoculate test sample aseptically.

Specimens types analyzed

Potable, waste water samples, air samples, food and dairy samples etc.

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Precautions to be taken

These microbial media are intended for the in-vitro use only. All the handling, experiments, storage, and discarding should be performed with the help of skilled and knowledgeable technicians and as per the established guidelines. The material should be disposed only after proper sterilization by autoclaving. Please go through the MSDS of the media to avoid any accidents or in emergency.

Performance and Evaluation

The expected performance of the medium is liable to use as per the direction on the label when stored at optimum conditions and within expiry date.

Quality Control

Appearance	Light beige colored free flowing, homogeneous powder
Reaction of 1.72 % solution	6.80 $\pm$ 0.2 at 25 °C
pH	6.60- 7.00
Color and clarity of ready medium	Light amber colored opalescent gel
Growth Promotion properties	Best at $\leq$ 100 CFU at 32-37 °C for 18-72 h
Indicative properties	Optimum at $\leq$ 100 CFU at 32-37 °C for 18-48 h
Negative control	Performed using sterile distilled water

Different Microbial Response: Prepare media as per the label directions. Inoculate and incubate at 35 $\pm$ 2°C for 18-24 hours.

Organism	ATCC	Inoculum (CFU)	Growth
<i>Escherichia coli</i>	8739	50-100	Luxuriant
<i>Klebsiella aerogenes</i>	13048	50-100	Luxuriant
<i>Pseudomonas aeruginosa</i>	27853	50-100	Luxuriant
<i>Bacillus spizizenii</i>	6633	50-100	Luxuriant
<i>Lactobacillus casei</i>	9595	50-100	Luxuriant
<i>Staphylococcus aureus</i>	25923	50-100	Luxuriant

This document has been produced electronically and it is valid without signature.

www.oxfordlabchem.com

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Storage and Shelf Life: The product is highly hygroscopic; keep the container tightly closed at all times and store it properly as per the conditions mentioned on the label. The declared expiry is valid only when stored as per the conditions mentioned on the label.

Note: Sterilize media immediately after reconstitution.

Disposal: To avoid the contamination or propagation of any hazardous microbes the used, unusable or modified preparation of this product must be disposed after autoclaving after completion of task.

Reference

1. Atlas, R. M. (2005). Handbook of media for environmental microbiology. CRC press.
2. Difco Manual (1998). 11th Edition. Difco Laboratories., Division of Becton Dickinson and Company, Sparks, Maryland, USA.

Disclaimer:

The information contained herein in good faith but makes no representations as to its comprehensiveness or accuracy. This document is intended only as a guide to the appropriate precautionary handling of the material by a properly trained person using this product. Individuals receiving the information must exercise their independent judgment in determining its appropriateness for a particular purpose.

Oxford Lab Fine Chem LLP makes no representations or warranties, either express or implied, including without limitation any warranties of merchantability, fitness for a particular purpose with respect to the information set forth herein or the product to which the information refers. Accordingly, Oxford Lab Fine Chem LLP will not be responsible for damages resulting from use of or reliance upon this information.